

BRAND NAME | Motrin

GENERIC NAME | Ibuprofen

**DIRECTIONS** Using the chart below, give the weight appropriate dosage listed every 6 hours as needed for a temperature greater than 101 degrees Fahrenheit or pain.

MORE INFORMATION KnowYourDose.org

WARNING DO NOT GIVE THIS MEDICATION TO A CHILD UNDER 6 MONTHS. Consult a physician if your child has any kidney problems.

REMEMBER Always read and follow the label on all medicines.

> Always put your medicine up and away and out of sight to avoid accidental ingestion by curious children.

Use the dosing device (dropper, syringe or cup) that comes with the product or pharmacist if you have any questions. a medication dosing syringe.

Never give adult medicines to children.

Whenever possible, use weight to determine your child's dose. Otherwise, use age.

Talk to your doctor, nurse or

Weight (pounds)	Infant Drops 50mg / 1.25 ml	Children's Liquid 100mg / 5ml
8-9	40 mg (1.0 ml)	
10-11	50 mg (1.25 ml)	
12-13	60 mg (1.5 ml)	
14-16	70 mg (1.75ml)	3.1 ml
17-18	80 mg (2.0 ml)	3.7 ml
19-22	90 mg (2.25ml)	4.2 ml
22-26	100 mg (2.5ml)	5 ml (1 tsp)
28-31	120 mg (3.0ml )	6 ml
32-35	140 mg (3.5ml)	7ml (11/2 tsp)
36-39	160 mg (4.0 ml)	8 ml
40-44		9 ml
45-48	_	10 ml (2 tsp)
49-53		11 ml
54-57		12 ml (2 1/2 tsp)
58-62	_	13 ml
63-66	_	14 ml
67-71		15 ml (3 tsp)
72-75		16 ml
76-80		17 ml (3 1/2 tsp)
81-84		18 ml
85-88		19 ml
89-93		20 ml (4 tsp)
94-99		20 ml (4 tsp)
≥ 100		20 ml (4 tsp)

## **BENADRYL**

Otherwise, use age.

syringe or cup) that comes with the product pharmacist if you have any questions.

Talk to your doctor, nurse or

<b>BRAND NAME</b>	Benadryl		
GENERIC NAME	Diphenhydramine		
DIRECTIONS	Using the chart below, give the weight appropriate dosage listed every 6 hours as needed for allergy symptoms or as recommended by your physician. This is the maximum recommended safe dosage based on your child's weight. If the listed dosage makes your child too sleepy, you may need to decrease the amount of medicine given in each dose.		
MORE INFORMATION	KnowYourDose.org		
WARNING	Consult a physician before giving this medication to children under <b>2 MONTHS</b> of age or if your child has any liver problems.		
REMEMBER	Always read and follow the label on all medicines.	Never give adult medicines to children.	
	Always put your medicine up and away and out of sight to avoid accidental	Whenever possible, use weight to determine your child's dose.	

ingestion by curious children.

Use the dosing device (dropper,

or a medication dosing syringe.

Weight (pounds)	Elixir (12.5mg / 5ml)	FastMelt Chewable (12.5mg tablets) Tablets, Capsules and Quick Dissolve Strips (25mg tablets or 25mg strips)	
8-9	1.25 ml (1/4 tsp)		
10-11	2 ml		
12-16	2.5 ml (1/2 tsp)		
17-21	3.75 ml (3/4 tsp)		
22-25	4 ml		
26-33	5 ml (1 tsp)	1 FastMelt	
33-39	6 ml	1 FastMelt	
40-45	7 ml (1 1/2 tsp)	1 1/2 FastMelt	
46-52	9 ml	1 1/2 FastMelt	
53-58	10 ml (2 tsp)	2 FastMelt	1 tab/cap/strip
59-65	11 ml	2 FastMelt	1 tab/cap/strip
66-72	12 ml (2 1/2 tsp)	2 1/2 FastMelt 1 1/2 tabs/caps/strips	
73-80	14 ml	2 1/2 FastMelt 1 1/2 tabs/caps/strips	
81-88	15 ml (3 tsp)	3 FastMelt	1 1/2 tabs/caps/strips
89-99	20 ml	4 FastMelt	2 tabs/caps/strips
100-109	20 ml	4 FastMelt	2 tabs/caps/strips
≥ 110	20 ml (4 tsp)	4 FastMelt	2 tabs/caps/strips

11955 Dallas Parkway, Suite 400 | Frisco, Texas 75033 214.396.5200 | PediatricPeople.com

## **TYLENOI**

<b>BRAND NAME</b>	Tylenol
GENERIC NAME	Acetaminophen
DIRECTIONS	Using the chart below, give the weight appropriate dosage listed every 4 hours as needed for a temperature greater than 101 degrees Fahrenheit or pain.
MORE INFORMATION	KnowYourDose.org

WARNING Consult a physician before giving this medication to children under **2 MONTHS** of age or if your child has any liver problems.

## REMEMBER | Always read and follow the label

on all medicines.

Always put your medicine up and away and out of sight to avoid accidental ingestion by curious children.

Use the dosing device (dropper, syringe or cup) that comes with the product or a medication dosing syringe.

Never give adult medicines to children.

Whenever possible, use weight to determine your child's dose. Otherwise, use age.

Talk to your doctor, nurse or pharmacist if you have any questions.

Weight (pounds)	Children's Liquid 160mg / 5ml	Chewable Tablets 80mg tablets	Junior Strength 160 mg tablets
6-7	1.4 ml		
8-9	1.8 ml		
10-11	2.3 ml		
12-13	2.8 ml	1 tab	
14-16	3.1 ml	1 tab	
17-18	3.7 ml	1 tab	
19-22	4.2 ml	1 tab	
22-26	5 ml (1 tsp)	2 tabs	1 tab/cap
28-31	6 ml	2 tabs	1 tab/cap
32-35	7 ml (1 1/2 tsp)	3 tabs	1 1/2 tabs/caps
36-39	8 ml	3 tabs	1 1/2 tabs/caps
40-44	9 ml	3 tabs	1 1/2 tabs/caps
45-48	10 ml (2 tsp)	4 tabs	2 tabs/caps
49-53	11 ml	4 tabs	2 tabs/caps
54-57	12 ml (2 1/2 tsp)	5 tabs	2 1/2 tabs/caps
58-62	13 ml	5 tabs	2 1/2 tabs/caps
63-66	14 ml	5 tabs	2 1/2 tabs/caps
67-71	15 ml (3 tsp)	6 tabs	3 tabs/caps
72-75	16 ml	6 tabs	3 tabs/caps
76-80	17 ml (3 1/2 tsp)	7 tabs	3 1/2 tabs/caps
81-84	18 ml	7 tabs	3 1/2 tabs/caps
85-90	19 ml	7 tabs	3 1/2 tabs/caps
91-93	20 ml (4 tsp)	8 tabs	4 tabs/caps
94-99	20 ml (4 tsp)	8 tabs	4 tabs/caps
≥ 100	20 ml (4 tsp)	8 tabs	4 tabs/caps